


# Effects of Vegetation Manipulation on Breeding Waterfowl in Prairie Wetlands —A Literature Review



19970319 019

**DISTRIBUTION STATEMENT A**

Approved for public release;  
Distribution Unlimited

UNITED STATES DEPARTMENT OF THE INTERIOR  
FISH AND WILDLIFE SERVICE  
*Fish and Wildlife Technical Report 3*

## **FISH AND WILDLIFE TECHNICAL REPORTS**

This publication series of the Fish and Wildlife Service (formerly the Bureau of Sport Fisheries and Wildlife) comprises reports of investigations related to fish or wildlife. Each is published as a separate paper, but for economy several may be issued in a single cover. The Service distributes a limited number of these reports for the use of Federal and State agencies and cooperators. See inside back cover for a list of recent issues.

Copies of this publication may be obtained from the Publications Unit, U.S. Fish and Wildlife Service, Washington, DC 20240, or may be purchased from the National Technical Information Service (NTIS), 5258 Port Royal Road, Springfield, VA 22161.

### **Library of Congress Cataloging-in-Publication Data**

Kantrud, Harold A.

Effects of vegetation manipulation on breeding waterfowl in prairie wetlands.

(Fish and wildlife technical report ; no. 3)

Bibliography: p.

Supt. of Docs. no.: I49.100:3

1. Waterfowl management—United States. 2. Wildlife habitat improvement—United States. I. Title. II. Series.  
SK361.K36 1985 639.9'7841 85-600326

# **Effects of Vegetation Manipulation on Breeding Waterfowl in Prairie Wetlands—A Literature Review**

By Harold A. Kantrud

UNITED STATES DEPARTMENT OF THE INTERIOR  
FISH AND WILDLIFE SERVICE  
*Fish and Wildlife Technical Report 3*  
Washington, D.C. • 1986

## Contents

	Page
Abstract .....	1
Problems .....	2
Management of Wetland Vegetation for Waterfowl .....	5
Burning .....	5
Grazing .....	7
Research Needs .....	10
References .....	11

# Effects of Vegetation Manipulation on Breeding Waterfowl in Prairie Wetlands—A Literature Review

By

Harold A. Kantrud

*U.S. Fish and Wildlife Service  
Northern Prairie Wildlife Research Center  
Jamestown, North Dakota 58401*

## Abstract

Literature on the effects of fire and grazing on the wetlands used by breeding prairie waterfowl is reviewed. Both dabbling and diving ducks and their broods prefer wetlands with openings in the marsh canopy. Decreased use is commonly associated with decreased habitat heterogeneity caused by tall, robust hydrophytes such as *Typha* spp. and other species adapted to form monotypes in the absence of disturbance. Nearly all previous studies indicate that reductions in height and density of tall, emergent hydrophytes by fire and grazing (unless very intensive) generally benefit breeding waterfowl. Such benefits are an increase in pair density, probably related to increased interspersions of cover and open water which decreases visibility among conspecific pairs, and improvements in their invertebrate food resources that result from increased habitat heterogeneity. Research needs are great because of the drastic changes that have accrued to prairie wetlands through fire suppression, cultivation, and other factors. The physical and biological environments preferred by species of breeding waterfowl during their seasonal and daily activities should be ascertained from future studies in wetland complexes that exist in the highest state of natural preservation. Long-term burning and grazing experiments should follow on specific vegetatively-degraded wetlands judged to be potentially important breeding areas. Seasonality, frequency, and intensity of treatments should be varied and combined and, in addition to measuring the response of the biotic community, the changes in the physical and chemical environment of the wetlands should be monitored to increase our knowledge of causative factors and possible predictive values.

The natural forces of climate, grazing, and fire were once the major factors controlling the abundance and species composition of vegetation in prairie wetlands. Breeding and migrant birds that used the wetlands evolved successfully under these influences, as evidenced by numerous accounts of large numbers and varieties of water birds present under pristine conditions.

Although wetland drainage has received the most publicity, other activities of European man had greatly changed prairie wetlands by the end of the 19th century. Domestic animals confined within fences sometimes grazed wetlands almost year-round. Wetlands near farmsteads often became highly eutrophic from barnyard and feedlot runoff water. Prairie fires, feared by both farmers and cattlemen, were suppressed whenever possible, which allowed

dead vegetation to accumulate in many wetlands. In agricultural areas, bottom soils of the shallowest, least permanent wetlands were regularly cultivated, even during wet years. In some years, wetlands with moderate water-retention ability could also be cultivated. During drought years, the bottom soils of more permanent water bodies were used to raise crops. The vegetation in all or part of some wetlands was mowed as often as possible for hay or bedding for livestock. Some wetlands were burned in the fall to reduce the amount of snow trapped in the basin or to discourage the spread of weeds; these wetlands could sometimes be cultivated the following spring.

In more recent times cultivation of steep slopes, use of row crops, and the practice of summer fallowing have caused much topsoil to move into the basins of countless prairie wetlands, further changing their

vegetative species composition and abundance. Dissolved salts and residues from agricultural chemicals probably have moved into many prairie wetlands. Irrigation practices have also altered the hydrology and vegetation of prairie wetlands. Finally, both herbage and woody vegetation have increased greatly in many wetlands in the eastern portion of the prairie pothole region. In this area, much livestock raising has been discontinued; thus, many formerly grazed or hayed wetlands that remain undrained now lie idle.

Although these land-use practices have undoubtedly affected the value of prairie wetlands to waterfowl and other birds, especially on privately owned lands, the effects have been only slightly less severe on many wetlands owned or managed by conservation agencies.

Research has emphasized bird habitat use, behavior, food habits, and recruitment of prairie-nesting waterfowl. Techniques used to manage upland nesting cover and, to a lesser degree, to control rates of hen and egg predation at upland sites are now fairly well developed. Yet little is known about practices that can rejuvenate vegetatively degraded prairie wetlands and restore their attractiveness to breeding waterfowl and other marsh and aquatic birds.

Wildlife problems associated with vegetation in wetlands and the response of wetland vegetation and animal populations to fire and grazing by domestic livestock are reviewed in this paper.

Weller (1978) stated that the theoretical basis for present marsh management techniques for wildlife is weak because of poor experimental design and inadequate evaluation of results; he encouraged the adoption of community-oriented management systems based on natural successional patterns that give benefits for a longer time and at lower cost than artificial systems. He identified burning and grazing as the systems most in need of study. Murkin (1979) also urged that the natural processes involved in the marsh cycle be studied; he stressed the importance of determining if semiopen marsh can be maintained in a productive state.

The natural fluctuation of water levels is probably the most important cause of vegetative change in prairie wetlands. Control of water levels has been extensively used to manipulate vegetation on many areas in the United States, but is not discussed here because such control is possible on only a small portion of the publicly owned wetlands in the prairie pothole region. Artificial management with costly herbicides, explosives, and sophisticated mechanical devices is also not considered here.

## Problems

Physical characteristics of wetland vegetation to aquatic birds was first given specific attention by Beecher (1942), who also found a correlation between numbers of plant communities and bird nests found in an Illinois wetland. Still, little is known of the relations between physical and biological factors of wetlands and their effect on waterfowl (Poston 1969a). Perhaps the most widely recognized evidence of the sensitivity of marsh birds to changes in the structure and density of wetland vegetation is the generally decreased use by waterfowl of wetlands covered by dense stands of tall emergent vegetation and their increased use of the open areas, the shallow water sparsely vegetated with short emergents, and exposed shorelines and mud flats. This phenomenon was evident in early studies and observations of adult breeding waterfowl (Pirnie 1935; Ward 1942; Mendall 1948; Bue et al. 1952; Dzubin 1955; Evans and Black 1956), and in later investigations (Munro 1963; Larsson 1969; Hopper 1972; March et al. 1973; Björk 1976; Piast 1982). Preferences of dabbling ducks (Anatinae) for wetlands with openings in the marsh canopy or for flooded emergent vegetation of a shorter type are well documented (Marshall 1952; Glover 1956; Johnsgard 1956; Smith 1968; Drewien and Springer 1969; Poston 1969b; Hines 1975; Weller 1975a; McEnroe 1976; Bishop et al. 1979). Diving ducks (Aythyinae), of course, show strong relations with open water areas (Hochbaum 1944; Siegfried 1976; Stoudt 1982).

Detailed studies have related the daily activity patterns of breeding waterfowl to the increased attractiveness of wetlands that contain an interspersed cover and open water. Such areas may provide better food resources according to Girard (1941), McDonald (1955), SOWLS (1955), Williams and Imber (1970), Courcelles and Bedard (1978), Beule (1979), Kaminski and Prince (1981b), and Murkin et al. (1982). Multiple regression analyses indicate that an increase in the ratio of open water to emergent vegetation may manifest itself in dabbling duck populations through better isolation of conspecific pairs, and may provide a cue to quality feeding habitat (Kaminski and Prince 1984).

Nest densities or hatching success may also be greater in broken versus solid stands of emergent marsh vegetation (McDonald 1955; Steel et al. 1956; Nelson and Dietz 1966; Mihelsons 1968; Ward 1968; Krapu and Duebbert 1974; Mednis 1974; Murkin

1979). The importance of openings or bare areas along shorelines for preening, resting, or waiting sites for adult waterfowl is also evident (McDonald 1955; Smith 1955; Sowls 1955; Sugden and Benson 1970; Williams and Imber 1970; Seymour 1974; Fog 1976). Partial destruction of *Typha* spp. stands by herbicides has resulted in a 300–400% increase in adult ducks per unit of shoreline (Keith 1961).

Waterfowl broods also prefer semiopen or open emergent vegetative cover, as shown by early observations and investigations (Bennett 1938; Wellein 1942; Stoudt 1944; Evans et al. 1952; Beard 1953; Berg 1956; Evans and Black 1956; Johnsgard 1956), and later studies by Keith (1961), Trauger (1967), Williams and Imber (1970), Bengtson (1971), Stoudt (1971), Sugden (1973), Whitman (1974, 1976), Mundinger (1975), Newton and Campbell (1975), Patterson (1976), and Wheeler and March (1979).

In the single instance where more broods were observed in closed stands of vegetation, Ignatoski (1966) postulated that nest success might have been higher there or that predation might have been greater in the more open areas. Studies showing that broods of dabbling ducks prefer semiopen marsh include those of Chura (1961), Perret (1962), Parnell and Quay (1965), Quame and Grewe (1970), Thompson (1974), Hines (1975), Courcelles and Bedard (1978), Mack and Flake (1980), Godin and Joyner (1981), Ringelman and Longcore (1982), Sjöberg and Danell (1982), and Talent et al. (1982). Similar results have been reported for diving ducks (Hochbaum 1944; Hildén 1964; Lokemoen 1966; Hilliard 1974; Stoudt 1982). Use of wetlands by broods increased as the number of vegetative types at the edge of the open water zone increased (Hopper 1972).

Other relations between breeding waterfowl and the physical features of their wetland habitat have been proposed. Openings in shoreline emergent vegetation may make nest sites on nearby uplands more easily accessible to hens (Mednis 1974; Mihelsons et al. 1974). Some studies indicate that waterfowl may be less susceptible to predation in more open situations (Furniss 1938; Beard 1953; Trauger 1967; Moller 1975) or that predator pressure may be buffered from waterfowl by the presence of other forms of prey in more open areas (Weller 1979). It has also been noted that a heavy buildup of marsh vegetation can make nesting islands accessible to predators (Mihelsons 1968). Rogers (1964) postulated that predation on lesser scaup (*Aythya affinis*) nests may have increased in situations where females were forced to walk, rather than swim, to their nests.

Other marsh-dwelling birds and mammals may benefit greatly from the presence of openings in marsh vegetation (Beard 1953; Seabloom 1958; Weller and Spatcher 1965; Willson 1966; Orians 1972; Vogl 1973; Gorenzel et al. 1982; Nudds 1982; Stenzel 1982). Such conditions may also result in avian communities of greater species diversity or richness (Weller and Spatcher 1965; Weller and Fredrickson 1973; Weller 1975a, 1978; Harris et al. 1981).

Biologists have often attributed decreased wetland use by aquatic birds to decreased habitat heterogeneity caused by disruption (usually a reduction) of natural ecological processes, resulting in domination by tall, robust hydrophytes in such genera as *Scirpus*, *Carex*, *Typha*, *Salix*, and *Phragmites* (Fig. 1). In the absence of these processes, autogenic successional processes tend to build dense stands of such hydrophytes in most wetlands (Walker 1959; Jahn and Moyle 1964; Whitman 1976). Prairie wetlands are particularly susceptible to the establishment of monotypes because of low gradient shorelines, small differences in soils or organic matter content within basins, and the ability of many species to survive under a wide range of water conditions (Hammond 1961; Walker and Coupland 1968).

*Typha* spp. has spread rapidly across a major portion of the prairie pothole region. For example, Metcalf (1931) and F. M. Uhler (personal communication) saw few *Typha*-dominated wetlands in North Dakota during 1917–25. Metcalf found only common cattail (*T. latifolia*) in North Dakota, and the species was listed only for “springy places and in the vicinity of freshwater lakes.” Since then, *T. angustifolia* and the extremely robust *T. “glauca”* (a presumed *T. latifolia* × *T. angustifolia* hybrid) have become dominant in thousands of prairie wetlands whose salinity ranges from fresh through slightly brackish (Stewart and Kantrud 1971).

*Typha* spp. is well-adapted to form monotypes (Linde et al. 1976). *Typha* seeds germinate under a wide range of water depths (Weller 1975b) and tolerate a wide range of soil types (Dean 1933). Older plants prevent competition from younger plants by autotoxicity (McNaughton 1968). Because shoot death in *Typha* spp. occurs late in the growing season, this plant’s competitive advantage over other species is probably enhanced (Davis and van der Valk 1978). A process of self-thinning allows individual *Typha* plants to grow large; decomposition of these large plants may take as long as 2 years (Mason and Bryant 1975). Mechanical control of *Typha* spp. is difficult and expensive (Nelson and Dietz 1966; Weller 1975b).

When tall, robust emergents such as *Typha* spp. dominate a wetland, drastic environmental changes occur. Less insolation of marsh soils and the water column caused by tall emergents and their litter may reduce or eliminate other species of plants in the understory (Bennett 1938; Buttery and Lambert 1965; Spence and Chrystal 1970; Vogl 1973) or lower productivity (Willson 1966). Submerged plants, in particular, require water of sufficient depth to reproduce (Anderson 1978; Courcelles and Bedard 1978), and the

buildup of litter and organic material from emergent species may reduce water depth or eliminate shallow water areas (Ward 1942; Walker 1959; Hammond 1961; Ward 1968; Beule 1979). Buildup of litter and the shading effect also may result in lower soil or water temperature and slower rates of plant decomposition (Willson 1966; Godshalk and Wetzel 1978). Various emergent species may decompose at different rates as the result of differences in species composition of macroinvertebrate populations (Danell and Sjöberg



**Fig. 1.** A prairie wetland unburned for more than 45 years; dense stands of *Phragmites australis* (foreground) and *Typha angustifolia* (background) lie offshore from the wet-meadow zone, which is dominated by a mature stand of *Salix amygdaloides*. The area has seldom been grazed by livestock. (Roberts Cty., South Dakota, 6 miles southwest of Rosholt; photo by H. A. Kantrud.)



1979). Thus the development of monotypic stands of emergents may effectively remove some of the variation in decomposer organisms that could act to maintain or increase vegetative heterogeneity.

## Management of Wetland Vegetation for Waterfowl

### *Burning*

Komarek (1976) stated that the fire ecology of wetlands was sorely in need of scientific study. General references (Kozlowski and Ahlgren 1974; Wright and Bailey 1982) indicate that burning of marsh vegetation releases nutrients, opens the canopy and detrital layer, and allows for increased insolation and resultant earlier warming of bottom soils. Biological productivity usually increases following fire, even though plant species composition may be altered. Little change in species composition usually occurs when perennial species with meristem at or below ground level are burned during their dormant period.

Fires were common in prairie wetland vegetation in the early 19th century, as evidenced by the accounts of early traders and travelers. For example, in 1803 Henry and Thompson (1965) recorded fire rushing through "low places covered with reeds and rushes." In 1858 or 1859, Boller (1972) saw a large conflagration spread for many miles after being set by American Indians in "dry rushes in the prairie bottoms." Denig (1961), writing about his experiences during 1833-54, noted that fire would sweep over ice through wetland vegetation.

### Impacts on Vegetation

Little is known about the environmental effects of fire in prairie wetlands. Much of the available information is obtained from general observations on wetlands where the fire frequency or season was unknown, or from fires set in a variety of vegetation types, usually on a nonexperimental basis. Hence, the results are often inconsistent and of minimal predictive value. Early studies by Lewis et al. (1928) indicated the changes in a few plant communities in central Alberta that could be expected in the presence or absence of a burning regime. Furniss (1938) noted that heavily lodged stands of *Typha latifolia* and *Scirpus validus* could be removed by fire in Saskatchewan

wetlands. Ward (1942) found that dense beds of *Phragmites australis* in Manitoba wetlands could be opened up by either spring or late summer burns, but that only late summer burning killed the "roots" (root-crowns). Grange (1949) observed that smartweeds (*Polygonum* spp.) disappeared because of competition from *Carex* spp., *Typha* spp., *Phragmites australis*, and various grasses. He stated that burning was probably the only effective method of stimulating smartweed growth in Wisconsin wetlands. Truax and Gunther (1951) used fall and winter burns to control undesirable vegetation at Horicon Marsh, Wisconsin. Annual burning was used to maintain the *Carex* spp. community in other Wisconsin wetlands (Thompson 1959). Tester and Marshall (1962) saw little change in species composition of wetland vegetation when Minnesota marshes containing low fuel volumes were burned. Smeins (1965) listed a few wetland plants found in North Dakota marshes with a history of burning. Schlichtemeier (1967) successfully removed dead stems of *P. australis* and *Scirpus* spp. with a winter burn, even though snow covered the bases of the plants. Vogl (1967) found burning generally favorable as a means of controlling woody plant invaders in Wisconsin wetlands. Smith (1969) stated that *Typha* spp. could quickly be destroyed by fire in Alberta wetlands. Beule (1979) concluded that burning was an ineffective control for *Typha* spp. in Wisconsin wetlands unless the peat layer was also burned. Gorenzel et al. (1981) found that fire failed to kill *Typha* spp. and *S. americanus* in a Colorado wetland. Thompson (1982) studied the seasonal effects of burning *P. australis* stands in a Manitoba wetland, and concluded that the changes in species composition and productivity produced by fall burns were intermediate between those produced by spring or summer burns.

In seasonal prairie wetlands, Stewart and Kantrud (1972) thought *Polygonum coccineum* increased after burning. However, Millar (1973) found no change in stands of *Carex atherodes*, *Scolochloa festucacea*, and *Eleocharis palustris* after repeated burning, which indicates these common plants of seasonal wetlands are extremely fire tolerant.

The aforementioned studies and observations do not provide managers with definitive, quantifiable information needed to formulate burn prescriptions in prairie wetlands. Research on prescribed burning for these wetlands for wildlife production was urged by Ward (1968) and Weller (1978), but to date almost all marsh burning for improvement of Waterfowl habitat has been done on migration or wintering areas (Sanderson and Bellrose 1969; Rutkowski 1978).

### Effects on Breeding Waterfowl

There is little substantive information about fire as it affects use of prairie wetlands by breeding waterfowl. Bennett (1938) and Furniss (1938) probably were the first to postulate that some benefits to breeding waterfowl could accrue from marsh burning. Bennett recommended shoreline burning to open dense stands of emergents to increase foods for blue-winged teal (*Anas discors*), whereas Furniss noted that crow predation on Saskatchewan duck nests may be less in marshes where heavily lodged, old-growth *Typha* spp. and *Scirpus* spp. stands were opened up or rejuvenated by fire. Cartwright (1942) suggested that burning dense, matted vegetation in Manitoba meadows would improve use by nesting ducks. Ward (1942) stated that burned openings in dense stands of *Phragmites australis* were heavily used by breeding ducks at the Delta Marsh, Manitoba. Grange (1949) noted that plants that produced seeds readily eaten by ducks were easily lost to competition from other plants and considered burning the only effective way to control plant

succession in Wisconsin wetlands. In South Dakota, Evans and Black (1956) noted that burning often improved use of wetlands by pairs of breeding waterfowl. Drewien and Springer (1969) observed that many burned wetlands lacked roosting cover in the spring, but that overall use of the wetlands by breeding pairs was not much affected.

Only a few experimental marsh burns have been conducted to study the effects on breeding waterfowl. Ducks showed increased use of winter-burned stands of *P. australis* and *Scirpus acutus* in wetlands in the Nebraska Sandhills (Schlichtemeier 1967). Ward (1968) found that in a Manitoba wetland fire opened up old stands of *P. australis* that formerly were almost devoid of duck nests and stimulated growth of *Scolochloa festuacea*, which supported highest duck nest densities. However, duck nest success was low the first year after a fire on low, *Poa pratensis* prairie in Iowa (Messinger 1974). A more detailed study was conducted by Björk (1976), who observed that in a Swedish wetland "severely damaged" by overgrowth of *Phragmites australis* and *Carex acuta*, burning and



Fig. 2. Prescribed spring burn being used to open a dense stand of *Phragmites australis* on the J. Clark Salyer National Wildlife Refuge. (Bottineau Cty., North Dakota, 4.5 miles southeast of Westhope; photo by R. Giese.)

mechanical methods of vegetation control resulted in much greater use of the area by breeding ducks, probably because of the presence of higher populations of chironomid insects. Prescribed burning of *P. australis* and *Typha* spp. during the dormant season is practiced on some National Wildlife Refuges (Fig. 2).

In the absence of water control, burning of vegetation in wetlands that naturally retain water only seasonally probably cannot be justified as a management practice for breeding waterfowl (Diirro 1982). Diirro found that increased early-season productivity of plants and invertebrates in basins burned the previous fall was offset by a general scarcity of water caused by the reduced snow-trapping ability of burned vegetation. In addition, snow accumulations tend to crush the softer vegetation found in seasonal wetlands, causing them to maintain an open or semiopen aspect during most springs. However, I believe that in pristine times vegetation in such wetlands would have burned more frequently than that found in more permanent wetlands. Long-term experiments on the effects of fire in the less permanent types of wetlands are needed.

### Grazing

Much more is known about the effects of grazing than of burning on wetland plant communities. Unless unusually severe, grazing results in greater plant species diversity and the development of more intricate patterns and sharper boundaries among plant communities (Bakker and Ruyter 1981). Livestock trampling may affect the height and density of wetland vegetation more than consumption (Hilliard 1974). Overgrazing may cause a decrease in primary production (Reimold et al. 1975), an increase in water turbidity (Logan 1975), and areas devoid of vegetation (Bassett 1980), as shown in Fig. 3. Adaptations of wetland plants to grazing include nodal rooting and unpalatability (Walker 1968; Walker and Coupland 1968). Marshes often show greater vegetative response to grazing than upland communities (Bassett 1980). Lists of species that increase or are tolerant of grazing in wetlands in or near the prairie pothole region have been published by Evans et al. (1952), Smith (1953), Harris (1954), Smeins (1965), Dix and Smeins



Fig. 3. Long-term overgrazing can destroy nearly all emergent vegetation in those shallow prairie wetlands having firm bottom soils. (Dickey Cty., North Dakota, 9 miles northwest of Forbes; photo by H. F. Duebbert.)

(1967), Walker and Coupland (1968), Stewart and Kantrud (1972), and Millar (1973).

#### Effects on Breeding Waterfowl

Most active management of waterfowl habitat through grazing by domestic livestock occurs on the wintering grounds, where the usual goal is to increase the availability of seeds of annual food plants (Griffith 1948; Neely 1967; Ermacoff 1968; Sanderson and Bellrose 1969). The effects of grazing on the quality of wetland habitat used by breeding waterfowl have received much attention during general investigations but little by experimental design. Early work by Bennett (1937) and Furniss (1938) on wetlands in Iowa and Alberta, respectively, indicated that overgrazing degraded habitat for ducks that nested along marsh borders or over water, but that nest density increased and egg predation by crows was less when densely vegetated shorelines were opened up by livestock. Sowls (1951) noted that ungrazed edges of wetlands attracted few breeding ducks and stated that ducks might increase if such areas were moderately grazed. Disturbed shorelines that would otherwise have supported dense growths of *Typha* spp. and *Scirpus* spp. probably supported higher densities of dabbling ducks in South Dakota stock ponds (Bue et al. 1952). Glover (1956) concluded that light to moderate grazing of shorelines after 1 July would not harm their value to waterfowl. Studies of man-made wetlands confirmed the deleterious effects of overgrazing on use of these wetlands by breeding ducks (Shearer 1960; Uhlig 1963). A study conducted in South Dakota (Sand Lake National Wildlife Refuge, unpublished annual reports, 1957–61) reported increased use of grazed shorelines by breeding ducks, especially green-winged teal (*Anas crecca*), northern pintail (*A. acuta*), and blue-winged teal. Salyer (1962) found that grazing was less harmful to breeding ducks when water areas increased in number and depth. Light grazing was recommended by Munro (1963) to help open *Typha* stands, thereby improving prairie wetlands for breeding waterfowl. Poston (1969b) postulated that light to moderate grazing would result in near optimum conditions for northern shoveler (*Anas clypeata*) on Alberta wetlands. The moderately grazed portion of the wetland shown in Fig. 4 represents an interspersion of cover and open water that is attractive to waterfowl.

Drewien and Springer (1969) were probably the first to report that breeding ducks move to roost in more heavily vegetated wetlands at night. These wetlands contained patchy, moderately dense stands of *Carex*

spp., *Polygonum coccineum*, *Scirpus* spp., *Scolochloa festuacea*, and *Typha* spp. The authors believed that lack of roosting cover did not limit densities of blue-winged teal on their South Dakota study area; sufficient roosting cover was always present because of other land-use practices, and even the overgrazed wetlands grew acceptable amounts of cover as the season progressed. During the day, the teal were found at higher densities on idle than on grazed wetlands; however, the authors inferred that this related to the proximity of upland nesting cover to the idle wetlands, rather than to differences among wetlands.

Kirsch (1969) found that pair use was lower on grazed North Dakota wetlands, but the differences between grazed versus idle wetlands were not significant. Cattle disturbance of duck nests was thought to be important during the study. Gjersing (1971) found high losses of duck nests due to livestock trampling around Montana reservoirs when the nests were within 7 yards of the shoreline. Winter grazing in Utah wetlands seemingly did not affect nesting dabbling ducks, but probably was harmful to divers (Hilliard 1974). In Denmark, Moller (1975) recommended grazing of wetlands during the nonbreeding season, and Fog (1976) believed that great portions of ungrazed marshes were lost to breeding ducks by the invasion of *Phragmites australis*. By using multiple regression analysis, McEnroe (1976) found that the percentage of shoreline grazed on natural wetlands was positively related to density indices (pairs per wetland) for the mallard (*Anas platyrhynchos*), gadwall (*A. strepera*), and blue-winged teal; however, intensity of grazing was negatively associated with densities of blue-winged teal and redhead (*Aythya americana*). A similar analysis of dabbling duck use of man-made wetlands in South Dakota showed species-specific preferences associated with differences in vegetation height, density, or diversity caused by grazing (Flake et al. 1977). The northern shoveler (*Anas clypeata*) made greatest use of pastured wetlands in England (Thomas 1980). The highest concentrations of breeding canvasback (*Aythya valisineria*) seen by Stoudt (1982) in a Manitoba study area were usually associated with pastured wetlands containing open or half-open surfaces and stands of *Scirpus acutus*.

#### Waterfowl Broods and Grazing

Relations between grazing and use of wetlands by waterfowl broods have also received some attention. Girard (1941) noted that broods would benefit if wetland shorelines in Montana were protected from

overgrazing. In Manitoba, the *Typha*-choked wetlands containing less than 10% open water received almost no use by duck broods (Evans et al. 1952). A history of light to moderate spring and fall grazing resulted in the open *Carex* spp. and *Scolochloa festuacea* habitat which was preferred by broods; ponds with broken stands of *Scirpus acutus* resulting from moderate to heavy grazing throughout the growing season were also extensively used. Broods were far more abundant on South Dakota livestock ponds with grassy shorelines than on those with mud shorelines created by overgrazing (Bue et al. 1952). Short emergent

growth in sparse stands caused by grazing also provided the best brood habitat on eastern Montana stock ponds (Smith 1953). Harris (1954) observed that heavily grazed areas dominated by *Scirpus* spp. and *Juncus* spp. received the most use by broods in Washington potholes. Overgrazing, especially of small wetlands, created unsuitable brood habitat in South Dakota (Evans and Black 1956). Keith (1961) noted a large increase in brood density after partial destruction of *Typha* spp. stands by herbicides on Alberta impoundments, and he recommended combining grazing and herbicide applications to rejuvenate marsh




Fig. 4. Moderate cattle grazing created the semiopen and more diverse plant community shown on the right side of the fence; the portion on the left remains idle. (Stutsman Cty., North Dakota, 2.5 miles east-southeast of Woodworth; photo by K. F. Higgins.)

edges for ducks. In Colorado, broods used lightly to moderately grazed wetlands far more than either moderately to heavily grazed wetlands or those that lay idle (Hopper 1972). Evans and Kerbs (1977) identified South Dakota impoundments having gently sloping shorelines and light use by livestock as water areas that would develop the natural vegetation structure preferred by broods. Nonetheless, if such wetlands contain appreciable amounts of *Typha* spp., its influence on brood use may be negative (Mack and Flake 1980). Canvasback broods in Manitoba reached highest densities in pastured wetlands containing less than 33% emergent vegetation (Stoudt 1982). Hudson (1983) found that duck brood densities were positively related to the amount of vegetation in Montana livestock ponds, but no ponds totally covered with emergents were censused, and all ponds were grazed.

#### Invertebrate Food and Grazing

Only a few investigators have mentioned response by invertebrate animals to herbage removal by livestock. Munro (1963) stated that grazing of *Typha*-dominated prairie wetlands would increase the planktonic algae that are the primary foods of invertebrates. Hopper (1972) believed that light to moderate grazing of flooded emergent vegetation would provide invertebrate foods for duck broods and also allow them easier access to shoreline feeding areas. Some very large invertebrates in salt marshes (such as crabs) may decrease under heavy grazing, but recovery is probably very rapid once grazing pressure is lessened (Reimold et al. 1975). Decreases of invertebrate animals caused by grazing of wetlands probably occur only when livestock are present in enough numbers to destroy aquatic vegetation (Logan 1975).

Invertebrates have been known to be important in the nutrition of breeding ducks since the early 1960's (Voights 1973). Indeed, invertebrate numbers and taxa may surpass all other measured physical and biological variables as indicators of wetland quality for breeding ducks (Joyner 1980), although it may be necessary to determine the behavior and distribution of the invertebrates in order to accurately predict which microhabitats will attract feeding ducks (Joyner 1982). Management of invertebrates for waterfowl was reviewed by Schroeder (1973) who recommended manipulation of cover through water control, grazing, and burning. He cautioned that such manipulations should favor a good interspersion of cover types without creating excessive siltation, undue fluctuations

of water levels during the nesting season, extensive reductions in plant abundance and diversity, and contamination of water supplies by toxic chemicals. When the invertebrate resources of prairie wetlands are manipulated by mechanical methods (Murkin 1979; Kaminski and Prince 1981a, 1981b), treatments are expensive and the response of breeding waterfowl is often small or of short duration.

### Research Needs

The prairie wetland complex has been severely degraded. Thus, it is too late to determine precisely the natural plant associations and structural types of vegetation historically preferred by waterfowl species during different phases of the breeding cycle. Other than drainage, cultivation, and siltation, the worst problem now is decreased waterfowl use caused by the regression of many of the semipermanent wetlands toward *Typha* spp. monotypes, and the encroachment of woody plants such as *Salix* spp. The problem is especially noticeable in the eastern portion of the prairie pothole region where livestock production has decreased, and many wetlands now lie idle. The problem is no less severe on much of the publicly owned land devoted to waterfowl production. In this instance, wetland managers seldom have the time, equipment, or manpower to properly manipulate vegetation on wetlands. More important, managers lack the information needed to obtain desirable, predictable results. However, much useful information can still result by studying existing wetland complexes that are subjected to various land uses or combinations of uses.

Much remains to be learned about the physical and biological environments preferred by species of breeding waterfowl during their seasonal and daily activities. This should be ascertained from existing wetland complexes that are in the highest state of natural preservation. Knowledge of the preferred feeding, nesting, loafing and roosting areas, and reactions and adaptations of the birds to climatic changes and predator pressure would aid in evaluating future experiments in marsh management.

Armed with a better understanding of the life history of individual species, burning and grazing treatments should be applied individually and in combination to selected prairie wetlands of various classes, salinity subclasses, vegetative types, and sizes that are most important to the commonest species of waterfowl.



Burning and grazing experiments should stress seasonality, frequency, and intensity, and the interactions of these variables should be measured. Effects of cover level (amount of emergent cover) should be separated, if possible, from the effects of cover configuration (size of clumps, shape of clumps, distances between clumps), as suggested by Murkin et al. (1982). The investigations should be long-term because of the drastic climatic fluctuations in the prairie pothole region. It would also be helpful if treated wetlands were dispersed over a broad geographical area to allow for differences in precipitation across the region. Studies should not be limited to the effects on waterfowl, but include the response of the vegetative community and the invertebrate food organisms of waterfowl. The response of other wetland vertebrates (primarily herbivores) to higher nutritive quality of burned wetland vegetation should also be measured as recommended by Smith et al. (1984).

Finally, changes to the physical and chemical environment should be monitored to increase our knowledge of causative factors involved in the biotic responses observed, and for the potential predictive values of abiotic factors in future marsh manipulations. Basic measurements should include winter snow accumulations, fluctuations in water depth and temperature among wetland vegetation zones, insolation, and standard water quality parameters.

## References

- Anderson, M. G. 1978. Distribution and production of sago pondweed (*Potamogeton pectinatus* L.) on a northern prairie marsh. *Ecology* 59:154-160.
- Bakker, J. P., and J. C. Ruyter. 1981. Effects of five years of grazing on a salt-marsh vegetation. *Vegetatio* 44:81-100.
- Bassett, P. A. 1980. Some effects of grazing on vegetation dynamics in the Camargue, France. *Vegetatio* 43:173-184.
- Beard, E. B. 1953. The importance of beaver in waterfowl management at the Seney National Wildlife Refuge. *J. Wildl. Manage.* 17:398-436.
- Beecher, W. J. 1942. Nesting birds and the vegetation substrate. Chicago Ornithological Society, Chicago, Illinois. 69 pp.
- Bengtson, S. A. 1971. Habitat selection of duck broods in Lake Myvatn area, north-east Iceland. *Ornis Scand.* 2:17-26.
- Bennett, L. J. 1937. Grazing in relation to the nesting of the bluewing teal. *Trans. N. Am. Wildl. Conf.* 2:393-397.
- Bennett, L. J. 1938. The blue-winged teal, its ecology and management. Collegiate Press, Ames, Iowa. 144 pp.
- Berg, P. F. 1956. A study of waterfowl broods in eastern Montana with special reference to movements and the relationship of reservoir fencing to production. *J. Wildl. Manage.* 20:253-262.
- Beule, J. D. 1979. Control and management of cattails in southeastern Wisconsin wetlands. *Wisc. Dep. Nat. Resour. Tech. Bull.* 112. 41 pp.
- Bishop, R. A., R. D. Andrews, and R. J. Bridges. 1979. Marsh management and its relationship to vegetation, waterfowl, and muskrats. *Proc. Iowa Acad. Sci.* 86:50-56.
- Björk, S. 1976. The restoration of degraded wetlands. Pages 349-354 in M. Smart, ed. *International Conference on the Conservation of Wetlands and Waterfowl*. International Waterfowl Research Bureau, Slimbridge, England.
- Boller, H. A. 1972. Among the Indians: four years on the upper Missouri, 1858-1862. M. M. Quaife, ed. University of Nebraska Press, Lincoln. 370 pp.
- Bue, I. G., L. Blankenship, and W. H. Marshall. 1952. The relationship of grazing practices to waterfowl breeding populations and production on stock ponds in western South Dakota. *Trans. N. Am. Wildl. Conf.* 17:396-414.
- Buttery, B. R., and J. M. Lambert. 1965. Competition between *Glyceria maxima* and *Phragmites communis* in the region of Surlingham Broad. I. The competition mechanism. *J. Ecol.* 53:163-181.
- Cartwright, B. W. 1942. Regulated burning as a marsh management technique. *Trans. N. Am. Wildl. Conf.* 7:257-263.
- Chura, N. J. 1961. Food availability and preferences of juvenile mallards. *Trans. N. Am. Wildl. Conf.* 26:121-134.
- Courcelles, R., and J. Bedard. 1978. Habitat selection by dabbling ducks in the Baie Noire marsh, southwestern Quebec. *Can. J. Zool.* 57:2230-2238.
- Danell, K., and K. Sjöberg. 1979. Decomposition of *Carex* and *Equisetum* in a northern Swedish lake: dry weight loss and colonization by macro-invertebrates. *J. Ecol.* 67:191-200.
- Davis, C. B., and A. G. van der Valk. 1978. Litter decomposition in prairie glacial marshes. Pages 99-130 in R. E. Good, D. F. Whigham, and R. L. Simpson, eds. *Freshwater wetlands*. Academic Press, New York.
- Dean, B. E. 1933. Effect of soil type and aeration upon root systems of certain aquatic plants. *Plant Physiol.* 8:203-222.
- Denig, E. T. 1961. Five Indian tribes of the upper Missouri. J. C. Ewers, ed. University of Oklahoma Press, Norman. 217 pp.
- Diirro, B. W. 1982. Effects of burning and mowing on seasonal whitetop ponds in southern Manitoba. M.S. Thesis. Iowa State University, Ames. 48 pp.
- Dix, R. L., and F. E. Smeins. 1967. The prairie, meadow, and marsh vegetation of Nelson County, North Dakota. *Can. J. Bot.* 45:21-58.
- Drewien, R. C., and P. F. Springer. 1969. Ecological relationships of breeding blue-winged teal to prairie potholes. Pages 102-115 in *Saskatoon Wetlands Seminar*. Can. Wildl. Serv., Rep. Ser. 6.
- Dzubin, A. 1955. Some evidences of home range in waterfowl. *Trans. N. Am. Wildl. Conf.* 20:278-298.

- Ermacoff, N. 1968. Marsh and habitat management practices at the Mendota Wildlife Area. Calif. Dep. Fish Game, Game Manage. Leaflet 12. 11 pp.
- Evans, C. D., and K. E. Black. 1956. Duck production studies on the prairie potholes of South Dakota. U.S. Fish Wildl. Serv., Spec. Sci. Rep.—Wildl. 32. 59 pp.
- Evans, C. D., A. S. Hawkins, and W. H. Marshall. 1952. Movements of waterfowl broods in Manitoba. U.S. Fish Wildl. Serv., Spec. Sci. Rep.—Wildl. 16. 47 pp.
- Evans, K. E., and R. R. Kerbs. 1977. Avian use of live-stock watering ponds in western South Dakota. U.S. For. Serv., Gen. Tech. Rep. RM-35. 11 pp.
- Flake, L. D., G. L. Petersen, and W. L. Tucker. 1977. Habitat relationships of breeding waterfowl on stock ponds in northwestern South Dakota. Proc. S. D. Acad. Sci. 56:135-151.
- Fog, J. 1976. Multi usage of waterfowl habitat. Pages 359-363 in M. Smart, ed. International Conference on the Conservation of Wetlands and Waterfowl. International Waterfowl Research Bureau, Slimbridge, England.
- Furniss, O. C. 1938. The 1937 waterfowl season in the Prince Albert District, central Saskatchewan. Wilson Bull. 50:17-27.
- Girard, G. L. 1941. The mallard: its management in western Montana. J. Wildl. Manage. 5:233-259.
- Gjersing, F. M. 1971. A study of waterfowl production on two rest rotation grazing units in north central Montana. M.S. Thesis. Montana State University, Bozeman. 42 pp.
- Glover, F. A. 1956. Nesting and production of the blue-winged teal (*Anas discors* Linnaeus) in northwest Iowa. J. Wildl. Manage. 20:28-46.
- Godin, P. R., and D. E. Joyner. 1981. Pond ecology and its influence on mallard use in Ontario, Canada. Wildfowl 32:28-34.
- Godshalk, G. L., and R. G. Wetzel. 1978. Decomposition in the littoral zone of lakes. Pages 131-143 in R. E. Good, D. F. Whigham, and R. L. Simpson, eds. Freshwater wetlands. Academic press, New York.
- Gorenzel, W. P., R. A. Ryder, and C. E. Braun. 1981. American coot response to habitat change on a Colorado marsh. Southwest. Nat. 26:59-65.
- Gorenzel, W. P., R. A. Ryder, and C. E. Braun. 1982. Reproduction and nest site characteristics of American coots at different altitudes in Colorado. Condor 84:59-65.
- Grange, W. B. 1949. The way to game abundance. Charles Scribner's Sons, New York. 365 pp.
- Griffith, R. 1948. Improving waterfowl habitat. Trans. N. Am. Wildl. Conf. 13:609-618.
- Hammond, M. C. 1961. Habitat improvement studies at Lower Souris National Wildlife Refuge—past, present and proposed. U.S. Fish Wildl. Serv. 8 pp. [mimeographed].
- Harris, H. J., G. Fewless, M. Milligan, and W. Johnson. 1981. Recovery processes and habitat quality in a freshwater coastal marsh following a natural disturbance. Pages 363-379 in B. Richardson, ed. Wetland values and management. Minnesota Water Planning Board, St. Paul.
- Harris, S. W. 1954. An ecological study of the waterfowl of the potholes area, Grant County, Washington. Am. Midl. Nat. 52:403-432.
- Henry, A., and D. Thompson. 1965. New light on the early history of the greater northwest: the manuscript journals of Alexander Henry and David Thompson 1799-1814. E. Coues, ed. Ross and Hains, Minneapolis. 1027 pp.
- Hildén, O. 1964. Ecology of duck populations in the island group of Valassaret, Gulf of Bothnia. Ann. Zool. Fenn. 1:153-277.
- Hilliard, M. A. 1974. The effects of regulated livestock grazing on waterfowl production. Utah Div. Wildl. Resour. Publ. 74-8. 96 pp.
- Hines, J. E. 1975. Gadwall nesting and brood ecology at Waterhen Marsh, Saskatchewan. M.S. Thesis. University of Regina, Regina, Saskatchewan. 218 pp.
- Hochbaum, H. A. 1944. The canvasback on a prairie marsh. The American Wildlife Institute, Washington, D.C. 201 pp.
- Hopper, R. M. 1972. Waterfowl use in relation to size and cost of potholes. J. Wildl. Manage. 36:459-468.
- Hudson, M. S. 1983. Waterfowl production on three age-classes of stock ponds in Montana. J. Wildl. Manage. 47:112-117.
- Ignatoski, F. J. 1966. The ecology of wild ducks in two central Michigan impoundment marshes. M.S. Thesis. Michigan State University, Lansing. 37 pp.
- Jahn, L. R., and J. B. Moyle. 1964. Plants on parade. Pages 293-304 in J. P. Linduska, ed. Waterfowl Tomorrow. U.S. Department of the Interior, Washington, D.C.
- Johnsgard, P. A. 1956. Effects of water fluctuation and vegetation change on bird populations, particularly waterfowl. Ecology 37:689-701.
- Joyner, D. E. 1980. Influence of invertebrates on pond selection by ducks in Ontario. J. Wildl. Manage. 44:700-705.
- Joyner, D. E. 1982. Abundance and availability of invertebrates in ponds in relation to dietary requirements of mallards and blue-winged teal at Luther Marsh. Ont. Field Biol. 36:19-34.
- Kaminski, R. M., and H. H. Prince. 1981a. Dabbling duck activity and foraging responses to aquatic macroinvertebrates. Auk 98:115-126.
- Kaminski, R. M., and H. H. Prince. 1981b. Dabbling duck and aquatic macroinvertebrate responses to manipulated wetland habitat. J. Wildl. Manage. 45:1-15.
- Kaminski, R. M., and H. H. Prince. 1984. Dabbling duck-habitat associations during spring in Delta Marsh, Manitoba. J. Wildl. Manage. 48:37-50.
- Keith, L. B. 1961. A study of waterfowl ecology on small impoundments in southeastern Alberta. Wildl. Monogr. 6. 88 pp.
- Kirsch, L. M. 1969. Waterfowl production in relation to grazing. J. Wildl. Manage. 33:821-828.
- Komarek, E. V., Sr. 1976. Fire ecology review. Proc. Tall Timbers Fire Ecol. Conf. 14:201-216.
- Kozlowski, T. T., and C. E. Ahlgren. 1974. Fire and ecosystems. Academic Press, New York. 542 pp.
- Krapu, G. L., and H. F. Duebbert. 1974. A biological survey of Kraft Slough. Prairie Nat. 6:33-55.
- Larsson, T. 1969. Land use and bird fauna on shores in southern Sweden. Oikos 20:136-155.
- Lewis, F. J., E. S. Dowding, and E. H. Moss. 1928. The vegetation of Alberta. II. The swamp, moor, and bog forest vegetation of central Alberta. J. Ecol. 16:19-70.



- Linde, A. F., T. Janisch, and D. Smith. 1976. Cattail: the significance of its growth, phenology and carbohydrate storage to its control and management. Wisc. Dep. Nat. Resour. Tech. Bull. 94. 27 pp.
- Logan, T. H. 1975. Characteristics of small impoundments in western Oklahoma, their value as waterfowl habitat and potential for management. M.S. Thesis. Oklahoma State University, Stillwater. 77 pp.
- Lokemoen, J. T. 1966. Breeding ecology of the redhead duck in western Montana. J. Wildl. Manage. 30:668-681.
- Mack, G. D., and L. D. Flake. 1980. Habitat relationships of waterfowl broods on South Dakota stock ponds. J. Wildl. Manage. 44:695-700.
- March, J. R., G. F. Martz, and R. A. Hunt. 1973. Breeding duck populations and habitat in Wisconsin. Wisc. Dep. Nat. Resour. Tech. Bull. 68. 37 pp.
- Marshall, W. H. 1952. Waterfowl of three prairie potholes. Flicker 24:60-68.
- Mason, C. F., and R. J. Bryant. 1975. Production, nutrient content, and decomposition of *Phragmites communis* Trin. and *Typha angustifolia* L. J. Ecol. 63:71-95.
- McDonald, M. E. 1955. Cause and effects of a die-off of emergent vegetation. J. Wildl. Manage. 19:24-35.
- McEnroe, M. R. 1976. Factors influencing habitat use by breeding waterfowl in South Dakota. M.S. Thesis. South Dakota State University, Brookings. 67 pp.
- McNaughton, S. J. 1968. Autotoxic feedback in relation to germination and seedling growth in *Typha latifolia*. Ecology 49:367-369.
- Mednis, A. A. 1974. Influence of vegetation on the species, composition and distribution of nesting ducks on the islands of Lake Engure. Pages 87-106 in H. A. Mihelsons and J. A. Viksne, eds. Ecology of waterfowl of Latvia [translated from Russian]. U.S. Dep. Commerce, Natl. Tech. Inf. Serv. TT71-53131.
- Mendall, H. L. 1948. Water levels and waterfowl management. Proc. 1948 Northeast. Game Conf. 86-88.
- Messinger, R. D. 1974. Effects of controlled burning on waterfowl nesting habitat in northwest Iowa. M.S. Thesis. Iowa State University, Ames. 49 pp.
- Metcalf, F. P. 1931. Wild-duck foods of North Dakota lakes. U.S. Dep. Agric. Tech. Bull. 221. 72 pp.
- Mihelsons, H. A. 1968. Methods of increasing the duck populations on lakes in Latvia. Pages 41-55 in Z. D. Spurin, chief ed. Birds of the Baltic region: ecology and migrations. Israel Program for Scientific Translations, Jerusalem.
- Mihelsons, H. A., G. T. Lejins, and A. A. Mednis. 1974. Nesting ducks of Lake Engure and their investigation. Pages 1-53 in H. A. Mihelsons and J. A. Viksne, eds. Ecology of waterfowl in Latvia [translated from Russian]. U.S. Dep. Commerce, Natl. Tech. Inf. Serv. TT71-53131.
- Millar, J. B. 1973. Vegetation changes in shallow marsh wetlands under improving moisture regime. Can. J. Bot. 51:1443-1457.
- Moller, H. S. 1975. Danish salt-marsh communities of breeding birds in relation to different types of management. Ornis Scand. 6:125-133.
- Mundinger, J. G. 1975. The influence of rest-rotation grazing management on waterfowl production on stock-water reservoirs in Phillips County, Montana. M.S. Thesis. Montana State University, Bozeman. 100 pp.
- Munro, D. A. 1963. Ducks and the Great Plains wetlands. Can. Audubon 25:105-111.
- Murkin, H. R. 1979. Response by waterfowl and black-birds to an experimentally manipulated cattail marsh. M.S. Thesis. McGill University, Montreal, Quebec. 97 pp.
- Murkin, H. R., R. M. Kaminski, and R. D. Titman. 1982. Responses by dabbling ducks and aquatic invertebrates to an experimentally manipulated cattail marsh. Can. J. Zool. 60:2324-2332.
- Neely, W. W. 1967. Planting, disking, mowing and grazing. Pages 212-221 in J. D. Newsom, ed. Proceedings of the Marsh and Estuary Management Symposium. Louisiana State University, Baton Rouge.
- Nelson, N. F., and R. H. Dietz. 1966. Cattail control methods in Utah. Utah Dep. Fish Game Publ. 66-2. 31 pp.
- Newton, I., and R. S. Campbell. 1975. Breeding of ducks at Loch Leven, Kinross. Wildfowl 26:83-103.
- Nudds, T. D. 1982. Ecological separation of grebes and coots: interference competition or microhabitat selection? Wilson Bull. 94:505-514.
- Orians, G. 1972. The adaptive significance of mating systems in the Icteridae. Proc. Fifteenth Int. Ornithol. Congr. 389-398.
- Parnell, J. F., and T. L. Quay. 1965. The populations, breeding biology, and environmental relations of the black duck, gadwall, and blue-winged teal at Pea and Bodie Islands, North Carolina. Proc. Southeast. Assoc. Game Fish Comm. 16:53-67.
- Patterson, J. H. 1976. The role of environmental heterogeneity in the regulation of duck populations. J. Wildl. Manage. 40:22-32.
- Perret, N. G. 1962. The spring and summer foods of the common mallard (*Anas platyrhynchos platyrhynchos* L.) in south central Manitoba. M.S. Thesis. University of British Columbia, Vancouver. 82 pp.
- Piest, L. A. 1982. Evaluation of waterfowl habitat improvements in the Apache-Sitgreaves National Forest, Arizona. M.S. Thesis. University of Arizona, Tucson. 87 pp.
- Pirnie, M. D. 1935. Michigan waterfowl management. Michigan Department of Conservation, Lansing. 328 pp.
- Poston, H. J. 1969a. Home range and breeding biology of the shoveler. M.S. Thesis. Utah State University, Logan. 86 pp.
- Poston, H. J. 1969b. Relationships between the shoveler and its breeding habitat at Strathmore, Alberta. Pages 132-137 in Saskatoon Wetlands Seminar. Can. Wildl. Serv., Rep. Ser. 6.
- Quame, O. A., and A. H. Grewe, Jr. 1970. Waterfowl production in a selected wetland management area. J. Minn. Acad. Sci. 36:89-92.
- Reimold, R. J., R. A. Linthurst, and P. L. Wolf. 1975. Effects of grazing on a salt marsh. Biol. Conserv. 8:105-125.
- Ringelman, J. K., and J. R. Longcore. 1982. Movements and wetland selection by brood-rearing black ducks. J. Wildl. Manage. 46:615-621.
- Rogers, J. P. 1964. Effect of drought on reproduction of the lesser scaup. J. Wildl. Manage. 28:213-222.
- Rutkowski, F. W. 1978. Bibliography on the effects of prescribed burning and mammal and waterfowl utilization of wetlands: annotated literature search with selected

- abstracts. Natl. Tech. Inf. Serv. (NTIS PB-286 117), Springfield, Virginia. 45 pp.
- Salyer, J. W. 1962. Effects of drought and land use on prairie nesting ducks. Trans. N. Am. Wildl. Nat. Resour. Conf. 27:69-79.
- Sanderson, G. C., and F. C. Bellrose. 1969. Wildlife habitat management of wetlands. An. Acad. Bras. Cienc. 41(Suppl.):153-204.
- Schlichtemeier, G. 1967. Marsh burning for waterfowl. Proc. Tall Timbers Fire Ecol. Conf. 6:41-46.
- Schroeder, L. D. 1973. A literature review on the role of invertebrates in waterfowl management. Colo. Div. Wildl. Spec. Rep. 29. 13 pp.
- Seabloom, R. W. 1958. A study of production, survival, and movements of the Great Plains muskrat (*Ondatra zibethica cinnamominus*) in North Dakota. M.S. Thesis. University of Minnesota, Minneapolis. 95 pp.
- Seymour, N. R. 1974. Territorial behavior of wild shovellers at Delta, Manitoba. Wildfowl 25:49-55.
- Shearer, L. A. 1960. Use of dugouts by breeding ducks. J. Wildl. Manage. 24:213-215.
- Siegfried, W. R. 1976. Segregation in feeding behavior of four diving ducks in southern Manitoba. Can. J. Zool. 54:730-736.
- Sjöberg, K., and K. Danell. 1982. Feeding activity of ducks in relation to diel emergence of chironomids. Can. J. Zool. 60:1383-1387.
- Smeins, F. E. 1965. The grassland and marshes of Nelson County, North Dakota. M.S. Thesis. University of Saskatchewan, Saskatoon. 82 pp.
- Smith, A. G. 1969. Waterfowl-habitat relationships on the Lousana, Alberta, waterfowl study area. Pages 116-122 in Saskatoon Wetlands Seminar. Can. Wildl. Serv., Rep. Ser. 6.
- Smith, L. M., J. A. Kadlec, and P. V. Fannesbeck. 1984. Effects of prescribed burning on nutritive quality of marsh plants in Utah. J. Wildl. Manage. 48:285-288.
- Smith, R. H. 1953. A study of waterfowl production on artificial reservoirs in eastern Montana. J. Wildl. Manage. 17:276-291.
- Smith, R. I. 1955. The breeding territory and its relationship to waterfowl productivity at Ogden Bay Bird Refuge. M.S. Thesis. Utah State Agriculture College, Logan. 46 pp.
- Smith, R. I. 1968. The social aspects of reproductive behavior in the pintail. Auk 85:381-396.
- Sowls, L. K. 1951. A study of the ecology and behavior of some surface-feeding ducks. Ph.D. Thesis. University of Wisconsin, Madison. 194 pp.
- Sowls, L. K. 1955. Prairie ducks. Stackpole Co., Harrisburg, Pennsylvania. 193 pp.
- Spence, D. H. N., and J. Chrystal. 1970. Photosynthesis and zonation of freshwater macrophytes. I. Depth distribution and shade tolerance. New Phytol. 69:205-215.
- Steel, P. E., P. D. Dalke, and E. G. Bizeau. 1956. Duck production at Grey's Lake, Idaho, 1949-1951. J. Wildl. Manage. 20:279-285.
- Stenzel, J. R. 1982. Ecology of breeding yellow rails at Seney National Wildlife Refuge. M.S. Thesis. Ohio State University, Columbus. 96 pp.
- Stewart, R. E., and H. A. Kantrud. 1971. Classification of natural ponds and lakes in the glaciated prairie region. U.S. Fish. Wildl. Serv., Resour. Publ. 92. 57 pp.
- Stewart, R. E., and H. A. Kantrud. 1972. Vegetation of prairie potholes, North Dakota, in relation to quality of water and other environmental factors. U.S. Geol. Surv., Prof. Pap. 585-D. 36 pp.
- Stoudt, J. H. 1944. Food preferences of mallards on the Chippewa National Forest, Minnesota. J. Wildl. Manage. 8:100-112.
- Stoudt, J. H. 1971. Ecological factors affecting waterfowl production in the Saskatchewan parklands. U.S. Fish. Wildl. Serv., Resour. Publ. 99. 58 pp.
- Stoudt, J. H. 1982. Habitat use and productivity of canvasbacks in southwestern Manitoba, 1961-72. U.S. Fish. Wildl. Serv., Spec. Sci. Rep.—Wildl. 248. 31 pp.
- Sugden, L. G. 1973. Feeding ecology of pintail, gadwall, American widgeon and lesser scaup ducklings. Can. Wildl. Serv., Rep. Ser. 24. 45 pp.
- Sugden, L. G., and D. A. Benson. 1970. An evaluation of loafing rafts for attracting ducks. J. Wildl. Manage. 34:340-343.
- Talent, L. G., G. L. Krapu, and R. L. Jarvis. 1982. Habitat use by mallard broods in south central North Dakota. J. Wildl. Manage. 46:629-635.
- Tester, J. R., and W. H. Marshall. 1962. Minnesota prairie management techniques and their wildlife implications. Trans. N. Am. Wildl. Nat. Resour. Conf. 27:267-287.
- Thomas, G. J. 1980. The ecology of breeding waterfowl at the Ouse Washes, England. Wildfowl 31:73-88.
- Thompson, D. J. 1982. Effects of fire on *Phragmites australis* (Cav.) Trin. ex Steudel and associated species at Delta Marsh, Manitoba. M.S. Thesis. University of Manitoba, Winnipeg. 199 pp.
- Thompson, D. Q. 1959. Biological investigation of the upper Fox River. Wisc. Conserv. Dep., Spec. Wildl. Rep. 2. 41 pp.
- Thompson, I. D. 1974. Movement and use of habitat by brood-rearing ducks at Lake St. Francis, Que. M.S. Thesis. York University, Toronto, Ontario. 123 pp.
- Trauger, D. L. 1967. Habitat factors influencing duck brood use of semi-permanent and permanent prairie potholes in North Dakota. M.S. Thesis. Iowa State University, Ames. 229 pp.
- Truax, W. C., and L. F. Gunther. 1951. The effectiveness of game management techniques employed on Horicon Marsh. Trans. N. Am. Wildl. Conf. 16:326-330.
- Uhlig, H. G. 1963. Use of Minnesota ponds and pits by waterfowl. Wilson Bull. 75:78-82.
- Vogl, R. J. 1967. Controlled burning for wildlife in Wisconsin. Proc. Tall Timbers Fire Ecol. Conf. 6:47-96.
- Vogl, R. J. 1973. Effects of fire on the plants and animals of a Florida wetland. Am. Midl. Nat. 89:334-347.
- Voights, D. K. 1973. Aquatic invertebrate abundance in relation to changing marsh conditions. Ph.D. Thesis. Iowa State University, Ames. 53 pp.
- Walker, B. H. 1968. Ecology of herbaceous wetland vegetation in the aspen grove and grassland regions of Saskatchewan. Ph.D. Thesis. University of Saskatchewan, Saskatoon. 180 pp.

- Walker, B. H., and R. T. Coupland. 1968. An analysis of vegetation-environment relationships in Saskatchewan sloughs. *Can. J. Bot.* 46:509-522.
- Walker, J. M. 1959. Vegetation studies on the Delta Marsh, Delta, Manitoba. M.S. Thesis. University of Manitoba, Winnipeg. 203 pp.
- Ward, E. 1942. Phragmites management. *Trans. N. Am. Wildl. Conf.* 7:294-298.
- Ward, P. 1968. Fire in relation to waterfowl habitat of the Delta marshes. *Proc. Tall Timbers Fire Ecol. Conf.* 8:254-267.
- Wellein, E. G. 1942. A waterfowl production study on the Chippewa National Forest. M.S. Thesis. University of Minnesota, Minneapolis. 71 pp.
- Weller, M. W. 1975a. Habitat selection by waterfowl of Argentine Isla Grande. *Wilson Bull.* 87:83-90.
- Weller, M. W. 1975b. Studies of cattail in relation to management for marsh wildlife. *Iowa State J. Res.* 49:383-412.
- Weller, M. W. 1978. Management of freshwater marshes for wildlife. Pages 267-284 in R. E. Good, D. F. Whigham, and R. L. Simpson, eds. *Freshwater wetlands*. Academic Press, New York.
- Weller, M. W. 1979. Small-mammal populations and experimental burning of Dewey's Pasture, northwest Iowa, 1970-1974. *Iowa State J. Res.* 53:325-332.
- Weller, M. W., and L. H. Fredrickson. 1973. Avian ecology of a managed glacial marsh. *Living Bird* 12:269-291.
- Weller, M. W., and C. E. Spatcher. 1965. Role of habitat in the distribution and abundance of marsh birds. *Iowa Agric. Home Econ. Exp. Sta., Spec. Rep.* 43. 31 pp.
- Wheeler, W. E., and J. R. March. 1979. Characteristics of scattered wetlands in relation to duck production in southeastern Wisconsin. *Wisc. Dep. Nat. Resour., Tech. Bull.* 116. 61 pp.
- Whitman, W. R. 1974. The response of macro-invertebrates to experimental marsh management. Ph.D. Thesis. University of Maine, Orono. 114 pp.
- Whitman, W. R. 1976. Artificial wetlands for waterfowl. Pages 336-344 in M. Smart, ed. *International Conference on the Conservation of Wetlands and Waterfowl*. International Waterfowl Research Bureau, Slimbridge, England.
- Williams, M., and M. J. Imber. 1970. The role of marginal vegetation in some waterfowl habitats. *N.Z. Ecol. Soc. Proc.* 17:126-128.
- Willson, M. F. 1966. Breeding ecology of the yellow-headed blackbird. *Ecol. Monogr.* 36:51-77.
- Wright, H. A., and A. W. Bailey. 1982. *Fire ecology*, United States and southern Canada. John Wiley & Sons, Inc. New York. 501 pp.

Kantrud, Harold A. **Effects of vegetation manipulation on breeding waterfowl in prairie wetlands—A Literature Review.** U.S. Fish Wildl. Serv., Fish Wildl. Tech. Rep. 3. 15 pp. 1986.

A literature review shows breeding ducks and their broods prefer wetlands with openings in the vegetative canopy. Dense growths of tall, robust hydrophytes decrease use of wetlands by breeding waterfowl. Burning and grazing generally increase habitat heterogeneity and degree of interspersed cover and open water. Such wetlands are more attractive to breeding ducks because of greater isolation of conspecific pairs and improvements in their invertebrate food resources. Research needs are great because prairie wetlands have been biologically degraded through fire suppression, cultivation and other factors. Research techniques are proposed.

**Keywords:** Waterfowl, wetlands, habitat, breeding, burning, grazing, foods, invertebrates, research.

Kantrud, Harold A. **Effects of vegetation manipulation on breeding waterfowl in prairie wetlands—A Literature Review.** U.S. Fish Wildl. Serv., Fish Wildl. Tech. Rep. 3. 15 pp. 1986.

A literature review shows breeding ducks and their broods prefer wetlands with openings in the vegetative canopy. Dense growths of tall, robust hydrophytes decrease use of wetlands by breeding waterfowl. Burning and grazing generally increase habitat heterogeneity and degree of interspersed cover and open water. Such wetlands are more attractive to breeding ducks because of greater isolation of conspecific pairs and improvements in their invertebrate food resources. Research needs are great because prairie wetlands have been biologically degraded through fire suppression, cultivation and other factors. Research techniques are proposed.

**Keywords:** Waterfowl, wetlands, habitat, breeding, burning, grazing, foods, invertebrates, research.

Kantrud, Harold A. **Effects of vegetation manipulation on breeding waterfowl in prairie wetlands—A Literature Review.** U.S. Fish Wildl. Serv., Fish Wildl. Tech. Rep. 3. 15 pp. 1986.

A literature review shows breeding ducks and their broods prefer wetlands with openings in the vegetative canopy. Dense growths of tall, robust hydrophytes decrease use of wetlands by breeding waterfowl. Burning and grazing generally increase habitat heterogeneity and degree of interspersed cover and open water. Such wetlands are more attractive to breeding ducks because of greater isolation of conspecific pairs and improvements in their invertebrate food resources. Research needs are great because prairie wetlands have been biologically degraded through fire suppression, cultivation and other factors. Research techniques are proposed.

**Keywords:** Waterfowl, wetlands, habitat, breeding, burning, grazing, foods, invertebrates, research.

Kantrud, Harold A. **Effects of vegetation manipulation on breeding waterfowl in prairie wetlands—A Literature Review.** U.S. Fish Wildl. Serv., Fish Wildl. Tech. Rep. 3. 15 pp. 1986.

A literature review shows breeding ducks and their broods prefer wetlands with openings in the vegetative canopy. Dense growths of tall, robust hydrophytes decrease use of wetlands by breeding waterfowl. Burning and grazing generally increase habitat heterogeneity and degree of interspersed cover and open water. Such wetlands are more attractive to breeding ducks because of greater isolation of conspecific pairs and improvements in their invertebrate food resources. Research needs are great because prairie wetlands have been biologically degraded through fire suppression, cultivation and other factors. Research techniques are proposed.

**Keywords:** Waterfowl, wetlands, habitat, breeding, burning, grazing, foods, invertebrates, research.

---

A list of current *Fish and Wildlife Technical Reports* follows.

1. Effects of Weather on Breeding Ducks in North Dakota, by Merrill C. Hammond and Douglas H. Johnson. 1984. 17 pp.
2. Lethal Dietary Toxicities of Environmental Contaminants and Pesticides to Coturnix, by Elwood F. Hill and Michael B. Camardese. 1986. In press.

As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.



UNITED STATES  
DEPARTMENT OF THE INTERIOR  
FISH AND WILDLIFE SERVICE  
OFFICE OF INFORMATION TRANSFER  
EDITORIAL SECTION  
1025 PENNOCK PLACE  
FORT COLLINS, COLORADO 80524

POSTAGE AND FEES PAID  
U.S. DEPARTMENT OF THE INTERIOR  
INT 423

**NOTE: Mailing lists are computerized. Please return address label with change of address.**